KIAS-P07026

Entropy of Schwarzschild Black Holes on DGP Brane

Ee Chang-Young*
Department of Physics, Sejong University, Seoul 143-747, Korea and
School of Physics, Korea Institute for Advanced Study, Seoul 130-722, Korea

Daeho Lee†
Department of Physics, Sejong University, Seoul 143-747, Korea

Myungseok Yoon‡
Department of Physics, Sejong University, Seoul, 143-747, Korea

(Dated: February 1, 2008)

arXiv:0706.2905v2 [gr-qc] 2 Nov 2007

Abstract

We study the entropy of Schwarzschild black holes on DGP brane. The radius of event horizon on DGP brane is obtained by a numerical method. It is smaller than that of Einstein gravity in the conventional branch, and is larger in the accelerated branch. However, the difference is very small. The entropy of the black hole is calculated by using the improved brick-wall method.

PACS numbers: 04.70.Dy,97.60.Lf

Keywords: Black Hole; Entropy

---

*Electronic address: cylee@sejong.ac.kr

†Electronic address: dhlee@sju.ac.kr

‡Electronic address: younms@sogang.ac.kr

After Bekenstein has suggested that the entropy of a black hole is proportional to the surface area at the event horizon [1], there have been several studies for the statistical origin of the entropy[2, 3, 4]. This can be understood by using a brick-wall method[5] or an *improved* brick-wall method[6, 7]. In the improved brick-wall method, a thin layer at the horizon is used instead of an infinite spherical box. As for a black hole on the Dvali-Gabadadze-Porrati(DGP) model[8], an exact expression of metric has not been obtained. In this paper, we first get the radius of event horizon numerically, then, based on this data about horizon, calculate the entropy of the black hole by using an improved brick-wall method.

The action in the DGP model is given by[8]

$$S = M_*^3 \int d^5x\sqrt{-g}R + M_P^2 \int d^4x\sqrt{-\tilde{g}}\tilde{R} \tag{1}$$

where the tilde denotes the quantities on the four-dimensional world-volume of the brane. A cross-over scale is defined by $r_c = m_c^{-1} = M_P^2/(2M_*^3)$. We consider a spherically symmetric and static black hole on the brane. Then, the five-dimensional line element can be written as

$$ds_{(5)}^2 = -e^{-\lambda}dt^2 + e^{\lambda}dr^2 + r^2 d\Omega^2 + \gamma dr dy + e^{\sigma}dy^2, \tag{2}$$

where λ, γ, and σ are functions only for r and y and the brane is located at $y = 0$. The equations of motion from the action (1) with Israel's junction conditions reduce to the equation for λ on the brane as follows[9]:

$$U_z^2 - 4(1+U)U_z - 8U(2+U) = 0, \tag{3}$$

where $z = \ln(r/r_0)$, $U = -2\exp(-2z)P_r/3m_c^2 r_0^2$, $P_r = dP/dr$, and $P(r) = r(1-\exp(-\lambda))|_{y=0}$ (r_0 being an arbitrary constant). Two solutions of Eq. (3), *regular* (or *conventional*) and *accelerated* branches, are given by[9]

$$k_1 r = R(U) = \left[-\frac{(1+3U+f)}{U^2(3+3U+\sqrt{3}f)^{2\sqrt{3}}(-5-3U+f)}\right]^{\frac{1}{8}}, \tag{4}$$

$$k_2 r = A(U) = \left[-\frac{(-5-3U+f)(-3-3U-\sqrt{3}f)^{2\sqrt{3}}}{(U+2)^2(1+3U+f)}\right]^{\frac{1}{8}}, \tag{5}$$

where k_1 and k_2 are constants of integration and $f = \sqrt{1+6U+3U^2}$. By applying boundary conditions by $P(0) = r_M$ and $P(\infty) = 0$ in the regular branch and $P(0) = r_M$ and $P(r) \approx m_c^2 r^3$ for large r in the accelerated branch, k_1 and k_2 are determined as $2(r_* k_1)^3 = c_1 \approx 0.43$



FIG. 1: Filled boxes and unfilled boxes are the ratios of r_H to r_M in the regular and the accelerated branches, respectively, where r_H is a horizon of a black hole on DGP brane and r_M is a horizon of a Schwarzschild black hole in Einstein gravity. It shows that r_H approaches r_M when r_c is much larger than r_M.

and $2(r_* k_2)^3 = c_2 \approx 4.41$, where $r_* = (r_M r_c^2)^{1/3}$[10]. From $P(r) = -\frac{3}{2} m_c^2 \int dr\ r^2 U(r)$, we obtain

$$P(r) = r_M \left[1 - \frac{1}{c_1} U R(U)^3 - \frac{1}{c_1}\int_U^\infty dU R(U)^3\right], \tag{6}$$

$$P(r) = r_M \left[1 - \frac{1}{c_2} U A(U)^3 + \frac{1}{c_2}\int_{-\infty}^U dU A(U)^3\right], \tag{7}$$

which correspond to the regular and accelerated branches, respectively.

Since $P(r_H) = r_H$ at the horizon, we can obtain $U_H \equiv U(r_H)$ from Eqs. (4) – (7). Substituting the corresponding U_H's into Eqs. (4) and (5), the horizons are calculated from $r_H/r_M = \left[2(r_c/r_M)^2/c_1\right]^{1/3} R(U_H)$ for the regular branch and $r_H/r_M = \left[2(r_c/r_M)^2/c_2\right]^{1/3} A(U_H)$ for the accelerated branch. The results for the regular and the accelerated branches are shown in Fig. 1. Note that the horizon can be written as $r_H = r_M + \xi$, where ξ should be very small because r_c/r_M is very large. This fact will be taken into account for calculating the entropy.

In order to find the entropy of a black hole, we would rewrite the line element (2) on the brane as $ds^2 = -f dt^2 + f^{-1} dr^2 + r^2 d\Omega^2$, where $d\Omega$ denotes a solid angle. The asymptotic behavior of the metric is obtained as

$$f(r) \approx 1 - \frac{r_M}{r} + b\alpha m_c^2 r^2 \left(\frac{r_*}{r}\right)^{2(\sqrt{3}-1)}, \quad \text{for } r \ll r_* \tag{8}$$

$$f(r) \approx 1 - \frac{\tilde{r}_M^2}{r^2} - b m_c^2 r^2, \qquad \text{for } r \gg r_* \tag{9}$$

where $\alpha \approx 0.84$ and $b = 1$ and $b = -1$ correspond to the conventional branch and the accelerated branch, respectively, [9, 10].

Since $a \equiv b\alpha m_c^2 r_*^{2(\sqrt{3}-1)}$ is very small, there exists a solution r_H such that $f(r_H) = 0$, where $r_H = r_M + \xi$ with $|\xi| \ll r_M$. The fact that ξ is very small compared to r_M appears in Fig. 1. From $f(r_H) = f(r_M + \xi) = 0$, we obtain $\xi \approx -a r_M^{5-2\sqrt{3}}$. Note that the radius of the event horizon is smaller than that of Einstein gravity in the conventional branch, and it is larger in the accelerated branch. Now, the Hawking temperature is given by $T = \beta^{-1} = \kappa/(2\pi) \approx \gamma/(4\pi r_H)$, where κ is a surface gravity of the black hole and $\gamma \equiv 1 - \xi/r_H + (4-2\sqrt{3}) a r_H^{4-2\sqrt{3}} \approx 1 - (5-2\sqrt{3})\xi/r_H$.

In order to calculate the entropy of a given system in the brick-wall method, we consider a quantum gas of scalar particles confined within a box near the horizon of a black hole and introduce a cut-off parameter[5]. The free scalar field is assumed to satisfy the Klein-Gordon equation, $(\Box + m^2)\Psi = 0$, with boundary conditions $\Psi(r_H + h) = \Psi(L) = 0$, where r_H is the horizon, m is the mass of a scalar field, h is an infinitesimal cut-off parameter, and $r_H + h$ and L represent the inner and the outer walls of a "spherical" box, respectively. Suppose that this system is in thermal equilibrium at a temperature T with an external reservoir. Using $\Psi = \exp(-i\omega t)\psi(r,\theta,\varphi)$, the Klein-Gordon equation is reduced to

$$\frac{\partial^2\psi}{\partial r^2} + \left(\frac{2}{r} + \frac{1}{f}\frac{\partial f}{\partial r}\right)\frac{\partial \psi}{\partial r} + \frac{1}{f}\left[\frac{\omega^2}{f} - m^2 + \frac{1}{r^2}\left(\frac{\partial^2}{\partial\theta^2} + \cot\theta\frac{\partial}{\partial\theta} + \csc^2\theta\frac{\partial^2}{\partial\varphi^2}\right)\right]\psi = 0. \quad (10)$$

By the WKB approximation, using $\psi \sim \exp[iS(r,\theta,\varphi)]$, we obtain $p_r^2 = f^{-1}[\omega^2/f - m^2 - p_\theta^2/r^2 - p_\varphi^2/(r^2\sin^2\theta)]$, where $p_r = \partial S/\partial r$, $p_\theta = \partial S/\partial\theta$, and $p_\varphi = \partial S/\partial\varphi$. Then, $p^2 = g^{rr}p_r^2 + g^{\theta\theta}p_\theta^2 + g^{\varphi\varphi}p_\varphi^2 = \omega^2/f - m^2$. The number of quantum states with energy not exceeding ω can be written as

$$n(\omega) = \frac{1}{(2\pi)^3}\int dr d\theta d\varphi dp_r dp_\theta dp_\varphi = \frac{2}{3\pi}\int dr\,\frac{r^2}{\sqrt{f}}\left(\frac{\omega^2}{f} - m^2\right)^{\frac{3}{2}}. \quad (11)$$

The free energy is given by

$$F = -\int_{m\sqrt{f}}^{\infty} d\omega\,\frac{n(\omega)}{e^{\beta\omega} - 1} = -\frac{2}{3\pi\beta^4}\int dr\frac{r^2}{f^2}\int_{x_0}^{\infty} dx\,\frac{(x^2 - x_0^2)^{3/2}}{e^x - 1}, \quad (12)$$

where $x = \beta\omega$ and $x_0 = \beta m\sqrt{f}$.

The degrees of freedom are dominant near horizon. Thus, we consider a thin layer with $L = r_H + h + \delta$, where δ is small. Since x_0 goes to zero near horizon, we obtain

$$F = -\frac{2}{3\pi\beta^4}\int dr\,\frac{r^2}{f^2}\int_0^{\infty} dx\,\frac{x^3}{e^x - 1} \approx -\frac{2\pi^3}{45\beta^4}\int_{r_H+h}^{r_H+h+\delta} dr\,\frac{r^2}{f^2}. \quad (13)$$

The metric near the horizon can be written as $f(r) \approx \gamma(1 - r_H/r)$. Therefore, the free energy can be calculated as

$$F \approx -\frac{2\pi^3 r_H^4}{45\beta^4\gamma^2}\frac{\delta}{h(h+\delta)}, \tag{14}$$

and the entropy becomes

$$S = \beta^2\frac{\partial F}{\partial\beta} \approx \frac{8\pi^2 r_H^4}{45\beta^2\gamma^2}\frac{\bar{\delta}^2}{\bar{h}^2(\bar{h}^2+\bar{\delta}^2)}, \tag{15}$$

where $\bar{h} \equiv \int_{r_H}^{r_H+h}\sqrt{g_{rr}}\,dr$, $\bar{\delta} \equiv \int_{r_H+h}^{r_H+h+\delta}\sqrt{g_{rr}}\,dr$. Note that $\bar{h}$ and $\bar{\delta}$ are the cut-off parameter and the thickness of the layer, respectively. Thus, we can write the entropy as

$$S \approx \frac{1}{4}\mathcal{A}\cdot\frac{1}{90\pi}\frac{\bar{\delta}^2}{\bar{h}^2(\bar{h}^2+\bar{\delta}^2)}, \tag{16}$$

where $\mathcal{A} = 4\pi r_H^2$. When the cut-off parameter $\bar{h}$ is chosen as $\bar{\delta}^2/[\bar{h}^2(\bar{h}^2+\bar{\delta}^2)] = 90\pi/G$, Eq. (16) agrees with the Bekenstein-Hawking entropy $S_{\rm BH} = \mathcal{A}/4G$[1, 2].

One may consider the cut-off introduced in the brick-wall method a bit ad hoc. In Ref.[11, 12], this point was criticized and the Pauli-Villars regularization scheme was used to replace the cut-off. In Refs.[13], the entropy was evaluated based on the notion of entanglement entropy. Below, we evaluate the entropy in the thin layer using the Pauli-Villars scheme and compare it with our result (16). Introducing the five regulator fields ($i = 1, \ldots, 5$) besides the original field ($i = 0$)[11], the total free energy becomes

$$\bar{F} \approx -\frac{2r_H^2}{3\pi\gamma^2}\sum_{i=0}^{5}\Delta_i\int_0^\infty\frac{d\omega}{e^{\beta\omega}-1}\int_{h'}^{L'}ds\frac{(\omega^2-\gamma m_i^2 s)^{3/2}}{s^2(1-s)^4}, \tag{17}$$

where $\Delta_0 = \Delta_3 = \Delta_4 = +1$ for the commuting fields and $\Delta_1 = \Delta_2 = \Delta_5 = -1$ for the anticommuting fields and the integration is taken for values where the square root is real. Setting $h' = 0$ and focusing only on the divergent contributions at the horizon, we obtain the entropy

$$\bar{S} \approx \frac{\pi r_H^3}{3\beta}\left(B + \frac{32\pi^2}{15\beta^2\gamma^2}A\right) = \frac{\mathcal{A}}{4}\frac{B\gamma}{12\pi} + \frac{A\gamma}{90}, \tag{18}$$

with the same A and B coefficients given in Ref.[11]. As it was discussed in Ref.[11], the second term including A corresponds to the 1-loop correction to the quadratic-curvatre terms of the gravitational acton, and is irrelevant to our case. The first term corresponds to the 1-loop renormalization of the Bekenstein-Hawking entropy $S = \mathcal{A}/4G_R$, where the renormalized Newton's constant G_R is given by $1/G_R = 1/G_B + B\gamma/12\pi$ with the bare Newton's constant G_B. Note that our result slightly differs from the result of Ref.[11] by a factor γ.

Acknowledgments

E. C-Y. and D. L. were supported by the Korea Research Foundation Grant funded by the Korean Government(MOEHRD), KRF-2006-312-C00498. E. C-Y. thanks KIAS for hospitality during the time that this work was done. M. Yoon was supported by the Korea Research Foundation Grant funded by the Korean Government (MOEHRD) (KRF-2005-037-C00017).

[1] J. D. Bekenstein, Lett. Nuovo. Cim. **4**, 737 (1972); Phys. Rev. D **7**, 2333 (1973); Phys. Rev. D **9**, 3292 (1974).

[2] S. W. Hawking, Commun. Math. Phys. **43**, 199 (1975).

[3] W. H. Zurek and K. S. Thorne, Phys. Rev. Lett. **54**, 2171 (1985).

[4] V. P. Frolov and I. Novikov, Phys. Rev. D **48**, 4545 (1993); V. P. Frolov and D. V. Fursaev, Phys. Rev. D **61**, 024007 (1999).

[5] G. 't Hooft, Nucl. Phys. B **256**, 727 (1985).

[6] F. He, Z. Zhao and S.-W. Kim, Phys. Rev. D **64**, 044025 (2001); W.-B. Liu and Z. Zhao, Chin. Phys. Lett. **18**, 310 (2001); Z. Zhou and W. Liu, Int. J. Mod. Phys. A **19**, 3005 (2004).

[7] W. Kim, Y-J. Park, E. J. Son, and M. S. Yoon, J. Kor. Phys. Soc. **49**, 15 (2006).

[8] G. Dvali, G. Gabadadze, and M. Porrati, Phys. Lett B **485**, 208 (2000).

[9] G. Gabadadze and A. Iglesias, Phys. Rev. D **72**, 084024 (2005).

[10] G. Gabadadze and A. Iglesias, Phys. Lett. B **632**, 617 (2006).

[11] J-G. Demers, R. Lafrance, and R. C. Myers, Phys. Rev. D **52**, 2245 (1995).

[12] S. P. Kim, S. K. Kim, K-S. Soh, and J. H. Yee, Phys. Rev. D **55**, 2159 (1997).

[13] T. Jacobson and R. Parentani, Phys. Rev. D **76**, 024006 (2007).